

Disclaimer

The following report(s) provides findings from an FDA-initiated query using Sentinel. While Sentinel queries may be undertaken to assess potential medical product safety risks, they may also be initiated for various other reasons. Some examples include determining a rate or count of an identified health outcome of interest, examining medical product use, exploring the feasibility of future, more detailed analyses within Sentinel, and seeking to better understand Sentinel capabilities.

Data obtained through Sentinel are intended to complement other types of evidence such as preclinical studies, clinical trials, postmarket studies, and adverse event reports, all of which are used by FDA to inform regulatory decisions regarding medical product safety. The information contained in this report is provided as part of FDA's commitment to place knowledge acquired from Sentinel in the public domain as soon as possible. Any public health actions taken by FDA regarding products involved in Sentinel queries will continue to be communicated through existing channels.

FDA wants to emphasize that the fact that FDA has initiated a query involving a medical product and is reporting findings related to that query does not mean that FDA is suggesting health care practitioners should change their prescribing practices for the medical product or that patients taking the medical product should stop using it. Patients who have questions about the use of an identified medical product should contact their health care practitioners.

The following report contains a description of the request, request specifications, and results from the modular program run(s).

If you are using a web page screen reader and are unable to access this document, please contact the Sentinel Operations Center for assistance at info@sentinelssystem.org.

Overview

Date Run: April 28, 2017

Request Description: The purpose of this report was to compare the frequency of diagnoses for gestational diabetes using International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) codes. ICD-10-CM code definitions were determined by mapping from ICD-9-CM code definitions using the Centers for Medicare and Medicaid Services (CMS) General Equivalence Mappings (GEMs). Simple forward mapping (SFM) and forward-backward Mapping (FBM) were used to map ICD-9-CM to ICD-10-CM codes.¹

Sentinel Modular Program Tool Used: Cohort Identification and Descriptive Analysis (CIDA) tool, version 3.3.6

Data Source: This request was run against the IBM® MarketScan® Commercial Claims and Encounters Database and Medicare Supplemental Database, which included 121 million members. Data from October 1, 2010 to March 31, 2016 were included in this report. The report includes three separate time periods: 1) October 1, 2010 to March 31, 2016, 2) January 1, 2015 to March 31, 2015, and 3) January 1, 2016 to March 31, 2016. See Appendix A for the dates of available data used in this report.

Study Design: We examined the incidence and prevalence of gestational diabetes across the ICD-9-CM era (October 2010 to September 2015) and ICD-10-CM era (October 2015 to March 2016) in the US. Incidence was additionally evaluated from January 2015 to March 2015 and January 2016 to March 2016. See Appendices B - D for specific codes.

Cohort Eligibility Criteria: Members included in the cohorts were required to be enrolled in plans with at least medical coverage. The following age groups were included in the cohorts: 2-17, 18-44, 45-64, 65+ years.

Incident Cohorts: Members included in the incident cohorts were required to be continuously enrolled in health plans with at least medical coverage for at least 183 days prior to gestational diabetes during which gaps in coverage of up to 45 days were allowed. Incident gestational diabetes was defined as no previous gestational diabetes in the 183 days preceding the index date with respect to ICD-9-CM and ICD-10-CM codes.

Prevalent Cohorts: There was no enrollment requirement for members in the prevalent cohorts. All qualifying diagnosis codes that occurred between October 1, 2010 and March 31, 2016 were included.

Limitations: Algorithms used to define outcomes are imperfect; thus, it is possible that there may be misclassification. Therefore, data should be interpreted with this limitation in mind. The MarketScan claims databases are based on a large convenience sample. Because the sample is not random, it may contain biases or fail to generalize well to other populations. Data come mostly from large employers; medium and small firms may be underrepresented.²

Please refer to Appendix E for detailed specifications of parameters used in the analyses for this request.

Notes: Please contact the Sentinel Operations Center Query Fulfillment Team (qf@sentinelssystem.org) for questions and to provide comments/suggestions for future enhancements to this document.

¹Fung, K. W., et al. (2016). "Preparing for the ICD-10-CM Transition: Automated Methods for Translating ICD Codes in Clinical Phenotype Definitions." EGEMS (Wash DC) 4(1): 1211.

²IBM Watson Health (2018). [online] ibm.com. Available at: <https://www.ibm.com/downloads/cas/OWZWJ0QO> [Accessed 01 Mar. 2019].

Table of Contents

<u>Glossary</u>	List of Terms Found in this Report and their Definitions
<u>Table 1</u>	Comparison of Incident Gestational Diabetes Diagnoses in the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Eras (January 1, 2015 - March 31, 2015 and January 1, 2016 - March 31, 2016)
<u>Figure 1</u>	Incidence of Gestational Diabetes Diagnoses per 1,000 Eligible Members from October 2010 - March 2016 by Code Type, 183-Day Washout
<u>Figure 2</u>	Prevalence of Gestational Diabetes Diagnoses per 1,000 Eligible Members from October 2010 - March 2016 by Code Type, 0-Day Washout
<u>Appendix A</u>	Dates Available for IBM® MarketScan® Commercial and Medicare Supplemental Databases
<u>Appendix B</u>	List of International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Codes Used to Define Gestational Diabetes
<u>Appendix C</u>	List of International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Codes Used to Define Gestational Diabetes, Simple Forward Mapping (SFM)
<u>Appendix D</u>	List of International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Codes Used to Define Gestational Diabetes, Forward-Backward Mapping (FBM)
<u>Appendix E</u>	Specifications for Parameters for this Request

Glossary of Terms for Analyses Using Cohort Identification and Descriptive Analysis (CIDA) Tool*

Amount Supplied - number of units (pills, tablets, vials) dispensed. Net amount per NDC per dispensing.

Blackout Period - number of days at the beginning of a treatment episode that events are to be ignored. If an event occurs during the blackout period, the episode is excluded.

Care Setting - type of medical encounter or facility where the exposure, event, or condition code was recorded. Possible care settings include: Inpatient Hospital Stay (IP), Non-Acute Institutional Stay (IS), Emergency Department (ED), Ambulatory Visit (AV), and Other Ambulatory Visit (OA). For laboratory results, possible care settings include: Emergency department (E), Home (H), Inpatient (I), Outpatient (O), or Unknown or Missing (U). Along with the Principal Diagnosis Indicator, forms the Care Setting/PDX parameter.

Ambulatory Visit (AV) - includes visits at outpatient clinics, same-day surgeries, urgent care visits, and other same-day ambulatory hospital encounters, but excludes emergency department encounters.

Emergency Department (ED) - includes ED encounters that become inpatient stays (in which case inpatient stays would be a separate encounter). Excludes urgent care visits.

Inpatient Hospital Stay (IP) - includes all inpatient stays, same-day hospital discharges, hospital transfers, and acute hospital care where the discharge is after the admission date.

Non-Acute Institutional Stay (IS) - includes hospice, skilled nursing facility (SNF), rehab center, nursing home, residential, overnight non-hospital dialysis and other non-hospital stays.

Other Ambulatory Visit (OA) - includes other non overnight AV encounters such as hospice visits, home health visits, skilled nursing facility visits, other non-hospital visits, as well as telemedicine, telephone and email consultations.

Cohort Definition (drug/exposure) - indicates how the cohort will be defined: (1): Cohort includes only the first valid treatment episode during the query period; (2): Cohort includes all valid treatment episodes during the query period; (3): Cohort includes all valid treatment episodes during the query period until an event occurs.

Days Supplied - number of days supplied for all dispensings in qualifying treatment episodes.

Eligible Members - number of members eligible for an incident treatment episode (defined by the drug/exposure and event washout periods) with drug and medical coverage during the query period.

Enrollment Gap - number of days allowed between two consecutive enrollment periods without breaking a "continuously enrolled" sequence.

Episodes - treatment episodes; length of episode is determined by days supplied in one dispensing or consecutive dispensings bridged by the episode gap.

Episode Gap - number of days allowed between two (or more) consecutive exposures (dispensings/procedures) to be considered the same treatment episode.

Event Deduplication - specifies how events are counted by the MP algorithm: (0): Counts all occurrences of an HOI during an exposure episode; (1): de-duplicates occurrences of the same HOI code and code type on the same day; (2): de-duplicates occurrences of the same HOI group on the same day (e.g., de-duplicates at the group level).

Exposure Episode Length - number of days after exposure initiation that is considered "exposed time."

Exposure Extension Period - number of days post treatment period in which the outcomes/events are counted for a treatment episode. Extensions days are added after any episode gaps have been bridged.

Lookback Period - number of days wherein a member is required to have evidence of pre-existing condition (diagnosis/procedure/drug dispensing).

Maximum Episode Duration - truncates exposure episodes after a requester-specified number of exposed days. Applied after any gaps are bridged and extension days added to the length of the exposure episode.

Member-Years - sum of all days of enrollment with medical and drug coverage in the query period preceded by an exposure washout period all divided by 365.25.

Minimum Days Supplied - specifies a minimum number of days in length of the days supplied for the episode to be considered.

Minimum Episode Duration - specifies a minimum number of days in length of the episode for it to be considered. Applied after any gaps are bridged and extension days added to the length of the exposure episode.

Monitoring Period - used to define time periods of interest for both sequential analysis and simple cohort characterization requests.

Principal Diagnosis (PDX) - diagnosis or condition established to be chiefly responsible for admission of the patient to the hospital. 'P' = principal diagnosis, 'S' = secondary diagnosis, 'X' = unspecified diagnosis, '.' = blank. Along with the Care Setting values, forms the Caresetting/PDX parameter.

Query Period - period in which the modular program looks for exposures and outcomes of interest.

Treatment Episode Truncation Indicator - indicates whether the exposure episode will be truncated at the occurrence of a requester-specified code.

Washout Period (drug/exposure) - number of days a user is required to have no evidence of prior exposure (drug dispensing/procedure) and continuous drug and medical coverage prior to an incident treatment episode.

Washout Period (event/outcome) - number of days a user is required to have no evidence of a prior event (procedure/diagnosis) and continuous drug and medical coverage prior to an incident treatment episode.

Years at Risk - number of days supplied plus any episode gaps and exposure extension periods all divided by 365.25.

*all terms may not be used in this report

Table 1. Comparison of Incident* Gestational Diabetes Diagnoses in the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) and International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Eras (January 1, 2015 - March 31, 2015 and January 1, 2016 - March 31, 2016)

	Members with Diagnosis	Eligible Members	Members with Diagnosis per 1,000 Eligible Members
Gestational Diabetes			
ICD-9-CM: January 1, 2015 - March 31, 2015	10,108	22,219,881	0.45
ICD-10-CM: Simple Forward Mapping: January 1, 2016 - March 31, 2016	9,473	22,695,591	0.42
ICD-10-CM: Forward-Backward Mapping: January 1, 2016 - March 31, 2016	10,331	22,694,158	0.46

* Incidence defined by 90 day washout

Figure 1. Incidence of Gestational Diabetes Diagnoses per 1,000 Eligible Members from October 2010 - March 2016 by Code Type, 183-Day Washout





Figure 2. Prevalence of Gestational Diabetes Diagnoses per 1,000 Eligible Members from October 2010 - March 2016 by Code Type, 0-Day Washout



Appendix A. Dates Available for IBM® MarketScan® Commercial and Medicare Supplemental Databases

Databases	Start Date	End Date
IBM MarketScan Commercial and Medicare Supplemental Databases ¹	1/1/2010	3/31/2016

¹ The IBM MarketScan Databases includes a sample of 121 million employees, dependents, and retirees in the United States with primary or Medicare supplemental coverage through privately insured fee-for-service, point-of-service, or capitated health plans. The IBM MarketScan claims databases are based on a large convenience sample. Because the sample is not random, it may contain biases or fail to generalize well to other populations. Data come mostly from large employers; medium and small firms may be underrepresented. For more information on the IBM MarketScan Databases, please review the White Paper here: <https://www.ibm.com/downloads/cas/OWZWJ0QO>

Appendix B. List of International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Codes Used to Define Gestational Diabetes

Code	Description	Code Type
648.8	Abnormal maternal glucose tolerance, complicating pregnancy, childbirth, or the puerperium	ICD-9-CM
648.80	Abnormal maternal glucose tolerance, complicating pregnancy, childbirth, or the puerperium, unspecified as to episode of care	ICD-9-CM
648.81	Abnormal maternal glucose tolerance, with delivery	ICD-9-CM
648.82	Abnormal maternal glucose tolerance, with delivery, with current postpartum complication	ICD-9-CM
648.83	Abnormal maternal glucose tolerance, antepartum	ICD-9-CM
648.84	Abnormal maternal glucose tolerance complicating pregnancy, childbirth, or the puerperium, postpartum condition or complication	ICD-9-CM

Appendix C. List of International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Codes Used to Define Gestational Diabetes, Simple Forward Mapping (SFM)

Code	Description	Code Type
O99.810	Abnormal glucose complicating pregnancy	ICD-10-CM
O24.419	Gestational diabetes mellitus in pregnancy, unspecified control	ICD-10-CM
O24.425	Gestational diabetes mellitus in childbirth, controlled by oral hypoglycemic drugs	ICD-10-CM
O24.429	Gestational diabetes mellitus in childbirth, unspecified control	ICD-10-CM
O99.814	Abnormal glucose complicating childbirth	ICD-10-CM
O24.435	Gestational diabetes mellitus in puerperium, controlled by oral hypoglycemic drugs	ICD-10-CM
O99.815	Abnormal glucose complicating the puerperium	ICD-10-CM
O24.415	Gestational diabetes mellitus in pregnancy, controlled by oral hypoglycemic drugs	ICD-10-CM
O24.439	Gestational diabetes mellitus in the puerperium, unspecified control	ICD-10-CM

Appendix D. List of International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) Codes Used to Define Gestational Diabetes, Forward-Backward Mapping (FBM)

Code	Description	Code Type
O99.810	Abnormal glucose complicating pregnancy	ICD-10-CM
O24.419	Gestational diabetes mellitus in pregnancy, unspecified control	ICD-10-CM
O24.425	Gestational diabetes mellitus in childbirth, controlled by oral hypoglycemic drugs	ICD-10-CM
O24.429	Gestational diabetes mellitus in childbirth, unspecified control	ICD-10-CM
O99.814	Abnormal glucose complicating childbirth	ICD-10-CM
O24.435	Gestational diabetes mellitus in puerperium, controlled by oral hypoglycemic drugs	ICD-10-CM
O99.815	Abnormal glucose complicating the puerperium	ICD-10-CM
O24.415	Gestational diabetes mellitus in pregnancy, controlled by oral hypoglycemic drugs	ICD-10-CM
O24.439	Gestational diabetes mellitus in the puerperium, unspecified control	ICD-10-CM
O24.410	Gestational diabetes mellitus in pregnancy, diet controlled	ICD-10-CM
O24.414	Gestational diabetes mellitus in pregnancy, insulin controlled	ICD-10-CM
O24.420	Gestational diabetes mellitus in childbirth, diet controlled	ICD-10-CM
O24.424	Gestational diabetes mellitus in childbirth, insulin controlled	ICD-10-CM
O24.430	Gestational diabetes mellitus in the puerperium, diet controlled	ICD-10-CM
O24.434	Gestational diabetes mellitus in the puerperium, insulin controlled	ICD-10-CM

Appendix E. Specifications for Parameters for this Request

Sentinel's Cohort Identification and Descriptive Analysis (CIDA) tool, version 3.3.6, was used to compare the frequency of diagnoses for gestational diabetes using International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-9-CM) versus ICD-10-CM codes.

Enrollment Gap: 45 days

Age Groups: 2-17, 18-44, 45-64, 65+ years

Enrollment Requirement: 183 days

Coverage Requirement: At least medical coverage

Scenario	Code Source	Query Start Date	Query End Date	Event				
				Event	Incident with respect to:	Washout (days)	Cohort Definition	Caresetting
1	ICD-9-CM Code Search	1/1/2015	3/31/2015	Gestational Diabetes	ICD-9-CM	90	First valid event only	Any
2	SFM	1/1/2016	3/31/2016	Gestational Diabetes	ICD-10-CM SFM	90	First valid event only	Any
3	FBM	1/1/2016	3/31/2016	Gestational Diabetes	ICD-10-CM FBM	90	First valid event only	Any
4	ICD-9-CM Code Search	10/1/2010	3/31/2016	Gestational Diabetes	ICD-9-CM or ICD-10-CM SFM	183	First valid event only	Any
6	SFM	10/1/2010	3/31/2016	Gestational Diabetes	ICD-9-CM or ICD-10-CM SFM	183	First valid event only	Any
7	FBM	10/1/2010	3/31/2016	Gestational Diabetes	ICD-9-CM or ICD-10-CM FBM	183	First valid event only	Any
8	ICD-9-CM Code Search	10/1/2010	3/31/2016	Gestational Diabetes	NA	0	All valid events	Any
9	SFM	10/1/2010	3/31/2016	Gestational Diabetes	NA	0	All valid events	Any
10	FBM	10/1/2010	3/31/2016	Gestational Diabetes	NA	0	All valid events	Any

ICD-9-CM and ICD-10-CM are provided by Optum360. ICD-10-CM codes were mapped from ICD-9-CM codes with the Centers for Medicare and Medicaid Services General Equivalence Mappings, using simple forward mapping (SFM) and forward-backward mapping (FBM).