

Disclaimer

The FDA chose a specific outcome algorithm that met its need for a given medical product-outcome assessment. The use of a specific outcome algorithm in a Sentinel assessment should not be interpreted as an endorsement from FDA to use the algorithm for all safety assessments. Investigators should always consider the objective, study design, analytic approach, and data source of a given medical product safety assessment when choosing the outcome algorithm. The suitability of an outcome algorithm may change when applied to different scenarios. For additional information, please refer to the [Best Practices for Conducting and Reporting Pharmacoepidemiologic Safety Studies Using Electronic Healthcare Data](#) guidance document provided by the FDA.

Overview

Title	Inpatient Depressive Disorder Algorithm Defined in "Neuropsychiatric Events following Montelukast Use: A Propensity Score Matched Analysis"
Request ID	cder_mpl2r_wp010
Description	<p>This report lists International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) diagnosis codes and algorithms used to define inpatient depressive disorder in this request.</p> <p>For additional information about the algorithm and how it was defined relative to the cohort and exposures of interest in the inferential analysis, see the analysis page here: https://www.sentinelinitiative.org/drugs/assessments/neuropsychiatric-events-following-montelukast-use-propensity-score-matched</p>
Outcome	Inpatient depressive disorder
Algorithm to Define Outcome	Evidence of an ICD-9-CM diagnosis code used to define depression in the inpatient care setting in the primary diagnosis position.
Query Period	January 1, 2000 - September 30, 2015
Request Send Date	July 17, 2019

Glossary

Care Setting - type of medical encounter or facility where the exposure, event, or condition code was recorded. Possible care settings include: Inpatient Hospital Stay (IP), Non-Acute Institutional Stay (IS), Emergency Department (ED), Ambulatory Visit (AV), and Other Ambulatory Visit (OA). For laboratory results, possible care settings include: Emergency department (E), Home (H), Inpatient (I), Outpatient (O), or Unknown or missing (U)

Outcome - outcome of interest (either primary or secondary)

Principal Diagnosis (PDX) - diagnosis or condition established to be chiefly responsible for admission of the patient to the hospital. 'P' = principal diagnosis, 'S' = secondary diagnosis, 'X' = unspecified diagnosis, '.' = blank. Along with the Care Setting values, forms the Caresetting/PDX parameter.

Query Period - period in which the modular program looks for exposures and outcomes of interest

Request Send Date - date the request was sent to Sentinel Data Partners

International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Diagnosis Codes Used to Define Inpatient Depressive Disorder in this Request

Codes	Description	Code Type	Code Category
311	Depressive disorder, not elsewhere classified	ICD-9-CM	Diagnosis
293.83	Mood disorder in conditions classified elsewhere	ICD-9-CM	Diagnosis
296.2	Major depressive disorder, single episode	ICD-9-CM	Diagnosis
296.20	Major depressive disorder, single episode, unspecified	ICD-9-CM	Diagnosis
296.21	Major depressive disorder, single episode, mild	ICD-9-CM	Diagnosis
296.22	Major depressive disorder, single episode, moderate	ICD-9-CM	Diagnosis
296.23	Major depressive disorder, single episode, severe, without mention of psychotic behavior	ICD-9-CM	Diagnosis
296.24	Major depressive disorder, single episode, severe, specified as with psychotic behavior	ICD-9-CM	Diagnosis
296.25	Major depressive disorder, single episode, in partial or unspecified remission	ICD-9-CM	Diagnosis
296.26	Major depressive disorder, single episode in full remission	ICD-9-CM	Diagnosis
296.3	Major depressive disorder, recurrent episode	ICD-9-CM	Diagnosis
296.30	Major depressive disorder, recurrent episode, unspecified	ICD-9-CM	Diagnosis
296.31	Major depressive disorder, recurrent episode, mild	ICD-9-CM	Diagnosis
296.32	Major depressive disorder, recurrent episode, moderate	ICD-9-CM	Diagnosis
296.33	Major depressive disorder, recurrent episode, severe, without mention of psychotic behavior	ICD-9-CM	Diagnosis
296.34	Major depressive disorder, recurrent episode, severe, specified as with psychotic behavior	ICD-9-CM	Diagnosis
296.35	Major depressive disorder, recurrent episode, in partial or unspecified remission	ICD-9-CM	Diagnosis
296.36	Major depressive disorder, recurrent episode, in full remission	ICD-9-CM	Diagnosis
296.5	Bipolar I disorder, most recent episode (or current), depressed	ICD-9-CM	Diagnosis
296.50	Bipolar I disorder, most recent episode (or current) depressed, unspecified	ICD-9-CM	Diagnosis
296.51	Bipolar I disorder, most recent episode (or current) depressed, mild	ICD-9-CM	Diagnosis
296.52	Bipolar I disorder, most recent episode (or current) depressed, moderate	ICD-9-CM	Diagnosis
296.53	Bipolar I disorder, most recent episode (or current) depressed, severe, without mention of psychotic behavior	ICD-9-CM	Diagnosis
296.54	Bipolar I disorder, most recent episode (or current) depressed, severe, specified as with psychotic behavior	ICD-9-CM	Diagnosis
296.55	Bipolar I disorder, most recent episode (or current) depressed, in partial or unspecified remission	ICD-9-CM	Diagnosis
296.56	Bipolar I disorder, most recent episode (or current) depressed, in full remission	ICD-9-CM	Diagnosis
296.90	Unspecified episodic mood disorder	ICD-9-CM	Diagnosis
298.0	Depressive type psychosis	ICD-9-CM	Diagnosis
300.4	Dysthymic disorder	ICD-9-CM	Diagnosis
301.12	Chronic depressive personality disorder	ICD-9-CM	Diagnosis
309.0	Adjustment disorder with depressed mood	ICD-9-CM	Diagnosis
309.1	Prolonged depressive reaction as adjustment reaction	ICD-9-CM	Diagnosis
309.28	Adjustment disorder with mixed anxiety and depressed mood	ICD-9-CM	Diagnosis